



Mark Scheme (Results)

Summer 2026

Pearson Edexcel International Advanced Level

In Statistics S1 (WST01) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN:

- bod – benefit of doubt
 - ft – follow through
 - the symbol $\checkmark$ will be used for correct ft
 - cao – correct answer only
 - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC – special case
 - oe – or equivalent (and appropriate)
 - d... or dep – dependent
 - indep – independent
 - dp – decimal places
 - sf – significant figures
 - * – The answer is printed on the paper or ag- answer given
 - $\square$ or d... – The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the

subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by 'MR' in the body of the script.
6. If a candidate makes more than one attempt at any question:
 - a) If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - b) If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme		Marks
1 (a)	[E(X) =] 3		B1
			(1)
(b)	$\left[E(X^2) = \sum x^2 P(X = x) = \right] 1^2 \times \frac{1}{5} + 2^2 \times \frac{1}{5} + 3^2 \times \frac{1}{5} + 4^2 \times \frac{1}{5} + 5^2 \times \frac{1}{5}$ or $\left[E(X^2) = \text{Var}(X) + E(X)^2 = \right] \frac{5^2 - 1}{12} + 3^2 [= 2 + 9]$		M1
	[E(X ²) =] 11		A1
			(2)
	Notes		Total 3
(a)	B1	cao Do not isw if they divide by 5 again after achieving 3	
(b)	M1	for a correct method to find $E(X^2)$ using $\sum x^2 P(X = x)$ Condone one omission. May be implied by 11 provided it is not an intermediate answer to an incorrect method to finding $E(X^2)$ Condone further work attempting to use 11 to find $\text{Var}(X)$ for this mark Alternatively score for use of $E(X^2) = \text{Var}(X) + E(X)^2$ May be implied by 2+'3' ²	
	A1	cao (correct answer with no incorrect working 2/2) do not isw if they divide by 5 again or they proceed to find $\text{Var}(X)$	

Question Number	Scheme		Marks
2 (a)	First Game Second Game Third Game		B1 B1 dB1 (3)
(b)	$0.9 \times 0.6 \times 0.3 = 0.162$		M1 A1 (2)
(c)	$0.9 \times 0.6 \times '0.7' + 0.9 \times '0.4' \times '0.6' + '0.1' \times '0.9' \times '0.6' [= '0.378' + '0.216' + '0.054']$ $= 0.648$		M1 dM1 A1 (3)
	Notes		Total 8
(a)	B1	for a complete structure with labels win and lose (Condone W and L for win and lose or other labels if clearly defined)	
	B1	for correct probabilities on branches for the first and second game	
	dB1	dep on both previous B marks it is for a fully correct tree diagram	
(b)	M1	for $0.9 \times 0.6 \times 0.3$ oe only May be implied by 0.16(2) but not if this is part of an incorrect method i.e. do not isw	
	A1	for 0.162 oe (e.g. $\frac{81}{500}$) (correct answer with no incorrect working 2/2)	
(c)	M1	for one correct product. You may ft on their tree diagram provided all are probabilities. They must be multiplying the 3 probabilities together. May be implied by one of 0.37(8) or 0.21(6) or 0.054 (allow rounded to 2sf or truncated)	
	dM1	dep on previous M. It is for all 3 correct products and attempting to add the three different products together. You may ft on their tree diagram of the correct structure provided all are probabilities. May be implied by $0.37(8) + 0.21(6) + 0.054$ (allow rounded to 2sf or truncated) Note if in (a) 1st B mark is B0 as they are missing the branches after WIN-LOSE then it is unlikely they will score this mark unless from a restart.	
	A1	for 0.648 oe (e.g. $\frac{81}{125}$) correct answer which has not clearly come from an incorrect method (3/3)	

Question Number	Scheme		Marks	
3 (a)	Width = 1.6[cm]		B1	
	[their 'h × w' =] awrt 21.0 or Area of 21[cm ² for a frequency of 14] or FD = 1.6 or k = 1.75 or 1.5		M1	
	Height = 13.125 (cm)		A1	
			(3)	
(b)	$[20.5] + \frac{32-21}{14} \times 5$ or $[20.5] + \frac{32.5-21}{14} \times 5$		M1	
	= awrt 24.4 or = awrt 24.6		A1	
			(2)	
(c) (i)	$\frac{1517.25}{64} = 23.707... = 23.71^*$		B1*	
	(ii)	$\sqrt{\frac{41368.3125}{64} - \left(\frac{1517.25}{64}\right)^2} = 9.1845....$ awrt 9.18	M1 A1	
			(3)	
(d)	Exact values are not known		B1	
			(1)	
(e)	Mean < Median so negative skew or Mean ≈ Median so no skew or $\frac{3("23.71" - "24.4"/"24.6")}{"9.1845"}$ so negative skew		B1ft	
			(1)	
	Notes		Total 10	
(a)	B1	for width = 1.6 oe ignore units		
	M1	for sight of 21 or 1.6 (provided this is not the width) or 1.75 or 1.5 which may be part of a calculation and may be unsimplified e.g. $21 = 14 \times 1.5$ so you may need to check this): Examples: $\left[\frac{\text{Height}}{\text{F.D.}} \Rightarrow \right] \frac{\text{Height}}{\frac{14}{5} (= 2.8)} = \frac{7.5}{'1.6'} \Rightarrow \text{Height} = \frac{21}{'1.6'} (\text{cm})$ Using areas of the bars $7.5 \times 3.2 = 24 \text{ cm}^2 = 16 \text{ photographers} \Rightarrow '1.6' \times h = \frac{14}{16} \times 24 \Rightarrow h = 13.125$ Using area proportional to frequency: Frequency ratio: $\frac{14}{16}$ Width ratio is $\frac{5}{10} = \frac{1}{2}$ $\Rightarrow \frac{\text{Height}}{7.5} \times \frac{1}{2} = \frac{14}{16} \Rightarrow \text{Height} = \left(\frac{14}{16} \times 2 \right) \times 7.5 [= 13.125]$ Direct ratio formula: $\frac{h_2}{h_1} = \frac{\frac{f_2}{w_2}}{\frac{f_1}{w_1}} \Rightarrow \frac{\text{Height}}{7.5} = \frac{\frac{14}{5}}{\frac{16}{10}} \Rightarrow \text{Height} = 7.5 \times \frac{14}{16} [= 13.125]$ May be implied by 13.1 or better		
		A1	13.125 oe (e.g. allow the exact fraction $\frac{105}{8}$ and allow 13.1 ignore units)	
		SC	width = awrt 1.42 and height = awrt 14.8 score B0M1A0	


(b)	M1	for a correct fraction including the sign to find the median using either n or $n + 1$ or allow the fraction including the sign if working from the upper limit e.g. $[25.5] - \frac{3}{14} \times 5$ Alternatively allow the equation: $\frac{32 - 21}{m - 20.5} = \frac{14}{5}$ oe. May be implied by $\frac{171}{7}$ or $\frac{689}{28}$ or a correct answer.
	A1	for awrt 24.4 or awrt 24.6 Do not accept a fraction for this mark.
(c)(i)	B1*	for a correct expression for the mean or 23.707... or better. Cannot be scored if only seen within the calculation for the standard deviation.
(ii)	M1	for a correct method to find the standard deviation including the square root i.e. $\sqrt{\frac{41368.3125}{64} - \mu^2}$ Follow through on their mean. May be implied by awrt 9.18 or $\sqrt{84.356..}$ May attempt $\sum fx^2$ using the table – look for an attempt with at least 3 correct products. Allow attempt at finding the s e.g. $\sqrt{\frac{41368.3125 - 64 \times '23.707...'^2}{63}} = 9.257...$ which may be implied by awrt 9.25 or awrt 9.26 or e.g. $\sqrt{85.6955...}$
	A1	awrt 9.18 or accept 9.185 or if using sample sd then awrt 9.26 or awrt 9.25 (correct answer scores 2/2)
(d)	B1	a correct reason given which explains that the exact values are not known. Do not allow references to the midpoints/exact values to score without further explanation that they are being used because the exact values are not known oe
(e)	B1ft	a correct comparison of the mean and median with a correct conclusion, ft their median or a correct calculation involving the given mean, their median and their standard deviation with a correct conclusion for their values. Note they may attempt to find the LQ and UQ for the data – they must be correct to at least 3sf if using n : $Q_1 = 17.4$, $Q_3 = 29.3$ if using $n + 1$ $Q_1 = 17.5$, $Q_3 = 29.5$ $Q_2 - Q_1 > Q_3 - Q_2 \Rightarrow$ negative skew but condone incorrect values in this attempt if they have also made a comparison of the mean with their median which is correct and the conclusion using the quartiles is consistent with this. Any comparison with the mode only is B0

Question Number	Scheme		Marks
4 (a)	$a + a^2 + 0.26 + 0.34 + 0.16 = 1$		M1
	$a^2 + a - 0.24 = 0 \Rightarrow (a - 0.2)(a + 1.2) = 0 \Rightarrow a = \dots$		dM1
	$a = 0.2$ as e.g. a cannot be negative *		A1*
			(3)
(b)	$E(X) = 1 \times 0.2 + 2 \times 0.2^2 + 3 \times 0.26 + 4 \times 0.34 + 5 \times 0.16 \left[= 3.22 = \frac{161}{50} \right]$		M1
	$E(2X + 4) = 2E(X) + 4 = 2 \times '3.22' + 4$		M1
	$= 10.44$ awrt 10.4		A1
			(3)
Alt(b)	$Y = 2X + 4$ so $y = 6, 8, 10, 12, 14$		M1
	$E(Y) = 6 \times 0.2 + 8 \times 0.2^2 + 10 \times 0.26 + 12 \times 0.34 + 14 \times 0.16 [= 10.44]$		M1
	$= 10.44$ awrt 10.4		A1
			(3)
(c)	$E(X^2) = 1^2 \times 0.2 + 2^2 \times 0.2^2 + 3^2 \times 0.26 + 4^2 \times 0.34 + 5^2 \times 0.16 \left[= 12.14 = \frac{607}{50} \right]$		M1
	$\text{Var}(X) = '12.14' - '3.22'^2 [= 1.7716]$		M1
	$\text{Var}(2X + 4) = 4\text{Var}(X) = 4 \times '1.7716'$		M1
	$= 7.0864$ awrt 7.09		A1
			(4)
Alt(c)	$Y = 2X + 4$ so $y^2 = 36, 64, 100, 144, 196$		M1
	$E(Y^2) = 36 \times 0.2 + 64 \times 0.2^2 + 100 \times 0.26 + 144 \times 0.34 + 196 \times 0.16 [= 116.08]$		M1
	$\text{Var}(Y) = '116.08' - '10.44'^2 [= 7.0864]$		M1
	$= 7.0864$ awrt 7.09		A1
			(4)
(d)	awrt “7.09”		B1ft
			(1)
Notes			Total 11
(a)	M1	for use of $\sum P(X = x) = 1$ which may be implied by a correct 3TQ	
	dM1	dep on 1st M for a 3TQ in a and attempts to solve the 3TQ by factorising, completing the square or quadratic formula proceeding to at least one value. If they state the roots without working (e.g. solving directly from a calculator) they must state both 0.2 and -1.2 oe. Correct roots from a correct 3TQ scores M1dM1 Condone when factorising if their factors are for the equivalent quadratic equation $\times 25$ e.g. $a^2 + a - 0.24 = 0 \Rightarrow (5a - 1)(5a + 6) = 0 \Rightarrow a = 0.2$	
	A1*	for achieving the given answer and a reason for selecting 0.2 following a correct quadratic. They either need to explain why they are rejecting the other root e.g. -1.2 is negative or probs are between 0 and 1 (or it is not a probability) or they need to explain why they are selecting 0.2 e.g. a must be greater than (or equal to) 0 Do not accept vague reasons e.g. -1.2 is not possible SC verification scores M1M0A0.	

Mark (b) and (c) together Look for values by the table for evidence of method or by the question		
(b)	M1	for use of $\sum xP(X = x)$ (with at least 3 correct products or terms with intention to add) but division by 5 is M0. May see 0.2+0.08+0.78+1.36+0.8 - cannot be implied by 3.22 alone. Look out for attempts at $2X$ which can score this mark. e.g. $2 \times 0.2 + 4 \times 0.04 + 6 \times 0.26 + 8 \times 0.34 + 10 \times 0.16$ (with at least 3 correct products or terms with intention to add) but division by 5 is M0
	M1	for use of $2E(X) + 4$ ft their $E(X)$ In the attempt at $2X$ for the first M they would need to add 4 for this mark
	A1	awrt 10.4 (provided all previous marks scored) Accept $\frac{261}{25}$
Alt(b)	M1	for $y = 6, 8, 10, 12, 14$ (May be seen in part (c)). They do not have to be evaluated. May be implied by the values 1.2, 0.32, 2.6, 4.08, 2.24
	M1	for use of $\sum yP(Y = y)$ (with at least 3 correct products or terms with intention to add) but division by 5 is M0. May see 1.2+0.32+2.6+4.08+2.24 which can imply the first two M marks - cannot be implied by awrt 10.4 alone
	A1	awrt 10.4 (provided all previous marks scored) Accept $\frac{261}{25}$
(c)	M1	for use of $\sum x^2P(X = x)$ (with at least 3 correct products or terms with intention to add) but division by 5 is M0. May see 0.2+0.16+2.34+5.44+4 Cannot be implied by 12.14 alone. Condone if labelled as $\text{Var}(X)$
	M1	for use of $E(X^2) - E(X)^2$ ft their values for $E(X^2)$ and $E(X)$
	M1	for use of $4\text{Var}(X)$ ft their positive $\text{Var}(X)$ – it cannot be for e.g. $4E(X^2)$ i.e. it must be a changed value from $E(X^2)$ even if it is labelled as $\text{Var}(X)$
	A1	awrt 7.09 (provided all previous marks scored)
Alt(c)	M1	for $y = 36, 64, 100, 144, 196$. They do not need to be evaluated. May see 7.2+2.56+26+48.96+31.36 Cannot be implied by 116.08 alone. Condone if labelled as $\text{Var}(X)$ If they attempt $2X$ the values 4, 16, 36, 64, 100 oe can score this mark (note they do not need to add 4 for the $\text{Var}(X)$)
	M1	for use of $\sum y^2P(Y = y)$ (with at least 3 correct products or terms with intention to add) or may attempt for just $2X$ (e.g. $4 \times 0.2 + 16 \times 0.04 + 36 \times 0.26 + 64 \times 0.34 + 100 \times 0.16$ oe) but division by 5 is M0
	M1	for use of $E(Y^2) - E(Y)^2$ ft their values for $E(Y^2)$ and $E(Y)$. May use the equivalent formula for $2X$
	A1	awrt 7.09 (provided all previous marks scored)
(d)	B1ft	awrt 7.09 or ft their positive answer in (c). Condone if their answer to (c) has subsequently been rounded to 3sf or better

Question Number	Scheme		Marks
5 (a)(i)	$\sum x^2 = 10(148.05 + 163.5^2) [= 268803] \quad *$ or $148.05 = \frac{(\sum x^2)}{10} - 163.5^2 \Rightarrow \frac{\sum x^2}{10} = 148.05 + 163.5^2 \Rightarrow \sum x^2 = 268803 \quad *$ or $148.05 = \frac{268803}{10} - 163.5^2 = 148.05 \Rightarrow \sum x^2 = 268803 \quad *$		B1*
	(ii)	$\sum y^2 = 10(205.25 + 61.5^2)$	M1
		$\sum y^2 = 39875$	A1
			(3)
	(b)(i)	$S_{yy} = 39875 - \frac{615^2}{10} [= 2052.5] \quad \text{or} \quad S_{yy} = 205.25 \times 10 [= 2052.5]$ or $S_{yy} = 39875 - 10 \times 61.5^2 [= 2052.5]$	
(ii)		$S_{xy} = 102230 - \frac{1635 \times 615}{10}$	M1
		$= 1677.5$	A1
			(3)
(c)		$\frac{'1677.5'}{\sqrt{1480.5 \times 2052.5}} = 0.9623...$ awrt 0.962	M1 A1
		(2)	
(d)	e.g. [Strong] positive correlation		B1ft
			(1)
(e)	e.g. The point (163.5, 61.5) is equal to $(\bar{x}, \bar{y})$		M1
	So PMCC will remain the same		A1
			(2)
Notes			Total 11
(a)(i)	B1*	for a correct expression for $\sum x^2$ on its own with sight of 10, 148.05 and 163.5 (or 26732.25) in their solution or if they have an equation with $\sum x^2$ as the only unknown, they need an intermediate stage of working before proceeding to 268803 Alternatively, show by verification that $\sum x^2 = 268803$ with a minimal conclusion. There should be no errors seen.	

(ii)	M1	for a correct method to find $\sum y^2$ May be implied by 39875
	A1	39875 (correct answer scores M1A1)
(b)(i)	B1*	for a correct expression for S_{yy}
(ii)	M1	for a correct method for S_{xy} May be implied by 1677.5
	A1	1677.5 oe (correct answer scores M1A1)
(c)	M1	for a correct method to find r using $S_{yy} = 2052$ and ft on their S_{xy} Does not need to be evaluated so do not be concerned by a value for r that is outside the valid range $[-1, 1]$. May be implied by awrt 0.96
	A1	awrt 0.962
(d)	B1ft	for indicating positive correlation provided $0 < r < 1$. Allow e.g. as x increases, y increases or positive linear relationship but positive relationship is B0 (too vague) ft on their r with a conclusion about correlation consistent with their value e.g. if $ r < 0.4$ allow no correlation oe May conclude negative correlation if $-1 < r < 0$
(e)	M1	for a correct reason about the point being equal to the means of x and y e.g. the values are the same as the means (of x and y)
	A1	for stay the same oe Note that M0A1 is not possible

Question Number	Scheme		Marks
6 (a)			B1
			(1)
(b)(i)	$[P(C) =] x + 0.2$		B1
(ii)	$[P(D C) =] \frac{0.2}{x + 0.2}$		B1
			(2)
(c)	$[P(C \cup D) =] x + y + 0.2$		B1
			(1)
(d)	$x + y + 0.44 = 1$ oe or $x + y + 0.2 = 0.76$ oe		M1
	$x + 0.2 = 3(y + 0.2)$ or $x + y + 0.2 = 0.6 + 4y \Rightarrow x - 3y = 0.4$		M1
	$y = 0.04$ and $x = 0.52$		A1A1
			(4)
(e)	$P(C) = 0.72, P(D) = 0.24, P(C \cap D) = 0.2$ or $P(C) = 0.72, P(C D) = 0.8333... $ or $P(D) = 0.24, P(D C) = 0.2777...$		M1
	$P(C) \times P(D) \neq P(C \cap D)$ or $P(C D) \neq P(C)$ or $P(D C) \neq P(D)$ So not independent		A1
			(2)
	Notes		Total 10
(a)	B1	for a correct shading – needs some indication of sections shaded / unshaded	
(b)(i)	B1	cao isw following a correct expression seen	
(ii)	B1	cao isw following a correct expression seen	
(c)	B1	cao Condone if they equate their expression to 0.76 only May be seen in (d) but must be an expression which is clearly labelled and not just embedded within an equation.	
(d)	M1	for either $x + y + 0.44 = 1$ oe or $x + y + 0.2 = 0.76$ oe May have already substituted in $x + 0.2 = 3(y + 0.2)$ e.g. $3y + 0.6 + y + 0.24 = 1$ which scores M1M1 Also allow if they state or find probabilities for x and y where $x + y = 0.56$ oe Note if the equations are seen elsewhere they must be used in this part to score.	
	M1	for $x + 0.2 = 3(y + 0.2)$ oe or $x + y + 0.2 = 0.6 + 4y$ oe may be seen within an equation	
	A1	for either $y = 0.04$ oe or $x = 0.52$ oe	
	A1	for both $y = 0.04$ oe and $x = 0.52$ oe	
	Note	Beware of e.g. $3(x + 0.2) = y + 0.2 \Rightarrow x = 0.04, y = 0.52$ which scores M1M0A0A0	
(e)	M1	for labelling all of the probabilities needed for a test of independence (probabilities must be correct or correct fit on their probabilities for x and y) Allow e.g. $0.2 + '0.52'$ as a probability. Must use C and D . The probabilities may be seen in a calculation which matches their stated test. Either $P(C)$, $P(D)$ and $P(C \cap D)$ or $P(D C)$ and $P(D)$ or $P(C D)$ and $P(C)$ (must all be probs)	
	A1	for stating the correct test with the correct probabilities $P(C) \times P(D) \neq P(C \cap D)$	

		or e.g. $P(D C) \neq P(D)$ and correct conclusion of not independent. If e.g. $P(C) \times P(D)$ is evaluated then it must be correct (allow to be rounded)	
Question Number	Scheme		Marks
7 (a)	$b = \frac{5780}{6948} = 0.83189...$		M1 A1
	$a = 62 - 'b' \times 55 = 16.2458...$		M1
	So $p = 16.2 + 0.832t$		A1
			(4)
(b)	Every extra mark in the theory test adds “0.832” marks to the practical test		B1ft
			(1)
(c)	$16.2 + 62 \times 0.832 [= 67.784]$ or $16.2 + 20 \times 0.832 [= 32.84]$		M1
(i) & (ii)	68 or awrt 67.8 33 or awrt 32.8/32.9		A1ft A1ft
			(3)
(d)	Student A is interpolation (and student B is extrapolation)		M1
	So the result for student A (is more reliable)		A1
			(2)
	Notes		Total 10
(a)	M1	for a correct method to find b Cannot be implied by awrt 0.83 but may be implied by a correct fraction. Condone $\frac{S_p}{S_t}$ to score this mark.	
	A1	awrt 0.83 oe (condone $\frac{1445}{1737}$ but not $\frac{5780}{6948}$) May be seen in further work / final equation	
	M1	for a correct method to find a Cannot be implied by awrt 16.2	
	A1	for $p =$ awrt $16.2 +$ awrt $0.832t$ (do not accept fractions for a and b for this mark) with all previous marks scored. Allow $p = 16.25$ Must be in terms of p and t Note that answer with no working scores 0 marks.	
	Note	May see attempts to use the straight-line equation e.g. $p - 62 = \frac{5780}{6948}(t - 55)$ which would score M1A0M1A0 until further work done to simplify the gradient and reach the required form. Send to review if unsure.	
(b)	B1ft	for a correct interpretation ft their gradient in context – must mention <u>theory (t)</u> , <u>practical (p)</u> and <u>marks</u> (condone scores). Must be clear that for each additional theory test mark, the practical mark increases by “0.832”	
(c) (i) (ii)	M1	for substitution of 62 or 20 into their regression line. May be implied by awrt 68 or awrt 33 or the answer using their regression line to 2sf. Allow truncated.	
	A1ft	for 68 or awrt 67.8 ft their regression line provided the answer is >0	
	A1ft	for 33 or awrt 32.8/32.9 ft their regression line provided the answer is >0	
	Note	If their regression line is not correct then you will need to check their answer Answers should be correct to 3sf but condone if they have given them to the nearest integer.	
(d)	M1	for a correct reason ‘ interpolation ’ oe on its own or allow reference to the marks. e.g. A as the value/mark in the theory test used is within the range. i.e. it has to be clear that it is the theory test mark (t) that is within the range of values (and not the practical mark) Mentioning 62 or 20 will imply they are referring to the theory test marks or referring to the range 21 to 97 (condone 23 to 80). Allow e.g. ‘62 because it is closer to the mean’ (it must be clear they are referring to the theory test mark)	

	A1	dep on M1 scored. It is for identifying student A (as more reliable) oe (e.g. allow 62 or 68 for A) Note that M0A1 is not possible	
Question Number	Scheme		Marks
8 (a)(i)	$\left[P(X < 440) = \right] P\left(Z < \frac{440 - 443}{3} \right)$		M1
	$\left[P(Z < -1) = 1 - 0.8413 \right] = 0.1587 \quad (\text{Calc } 0.158655\dots)$ awrt 0.159		A1
	$\left[P(441.5 < X < 447.5) = \right] P\left(\frac{441.5 - 443}{3} < Z < \frac{447.5 - 443}{3} \right)$		M1
	$\left[P(-0.5 < Z < 1.5) = 0.9332 - 0.3085 \right] = 0.6247 \quad (\text{Calc } 0.62465\dots)$ awrt 0.625		A1
			(4)
(b)	$\frac{443 + k - 443}{3} = 2.3263 \quad \text{or} \quad \frac{443 - k - 443}{3} = -2.3263$		M1 B1
	$k = 6.9789$ awrt 6.98		A1
			(3)
(c)	$P(Y < 338.1) = 0.025 \Rightarrow \frac{338.1 - \mu}{\sigma} = -1.96$ or $P(Y > 344) = 0.30 \Rightarrow \frac{344 - \mu}{\sigma} = 0.5244$		M1 A1A1
	$\left[5.9 = 2.4844\sigma \Rightarrow \right] \sigma = \dots, \mu = \dots$		dM1
	$\sigma = 2.3748\dots \quad \text{and} \quad \mu = 342.7546\dots$ awrt 2.37 and awrt 343		A1
			(5)
Notes			Total 12
	Note	Standardising using $\sigma = 9$ scores maximum (a)(i) M0A0 (ii) M0A0 (b)M0B1A0	
(a)(i)	M1	for standardising using 443 and 3 oe (implied by ± 1)	
	A1	awrt 0.159 and evidence of standardisation is seen (answer only is 0/2)	
(ii)	M1	for standardising both parts using 443 and 3 oe (implied by ± 0.5 and ± 1.5). It cannot be implied by $0.9332 - 0.3085$	
	A1	awrt 0.625 and evidence of standardisation is seen (answer only is 0/2)	
(b)	M1	for achieving an expression of $\pm \frac{k}{3}$ which may be unsimplified. May be seen in an equation or within a probability statement. Look out for use of e.g. $x = 443 + k$ and standardise $\frac{x - 443}{3}$ which would still score this mark. Condone if they use k for 'x' provided evidence of substituting back correctly in later work.	
	B1	for sight of ± 2.3263 or better	
	A1	awrt 6.98 (correct answer with no incorrect method seen scores 3/3) Allow 6.96 for use of $z = \pm 2.32$ and allow 6.99 for use of $z = \pm 2.33$	

		Note that M1B0A1 is possible
Alt(b)	M1	Sets $443 + k = a$ where $448 < a < 452$ or sets $443 - k = b$ where $434 < b < 438$
	B1	for sight of either 449.97(90438...) so awrt 449.98 or 436.02(0956...) so awrt 436.02
	A1	awrt 6.98 (correct answer with no incorrect method seen scores 3/3)
(c)	M1	for standardising with 338.1 and σ and setting = to a z value where $1.9 < z < 2$ or for standardising with 344 and setting = to a z value where $0.5 < z < 0.6$
	A1	for one correct equation with z value 3sf or better with compatible signs
	A1	for both equations correct with z values 3sf or better with compatible signs
	dM1	dep on the previous M. It is for solving simultaneously their two equations in μ and σ and proceeding to values for both μ and σ . Condone if σ is negative for this mark. You do not need to check this on your calculator.
	A1	dep on all previous marks scored it is for awrt 2.37 (also accept 2.375) and awrt 343 and both z values seen are 3sf or better